

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C07K 14/47, A61K 38/17, G01N 33/68</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/05254</b> <b>(43) International Publication Date:</b> 3 February 2000 (03.02.00)
<b>(21) International Application Number:</b> PCT/EP99/05050 <b>(22) International Filing Date:</b> 16 July 1999 (16.07.99) <b>(30) Priority Data:</b> 98202470.5                      23 July 1998 (23.07.98)                      EP <b>(71) Applicant (for all designated States except US):</b> AKZO NOBEL N.V. [NL/NL]; Velperweg 76, NL-6824 BM Arnhem (NL). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> VERHEIJDEN, Gijsbertus, Franciscus, Maria [NL/NL]; Heischouw 7, NL-5345 XT Oss (NL). BOOTS, Anna, Maria, Helena [NL/NL]; Verlengde Torenstraat 10, NL-5366 AV Megen (NL). <b>(74) Agent:</b> HERMANS, F., G., M.; P.O. Box 20, NL-5340 BH Oss (NL).		<b>(81) Designated States:</b> AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>

**(54) Title:** NOVEL PEPTIDES FOR USE IN IMMUNOTHERAPY OF AUTOIMMUNE DISEASES**(57) Abstract**

The invention relates to the use of novel peptides in a peptide induced tolerance therapy to prevent autoimmune disorders and in particular their use in treatment of chronic destruction of articular cartilage. The invention furthermore embraces pharmaceutical compositions comprising said peptides and a diagnostic method for the detection of autoreactive T cells in a test sample.

**BEST AVAILABLE COPY**

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark						

## NOVEL PEPTIDES FOR USE IN IMMUNOTHERAPY OF AUTOIMMUNE DISEASES

5       The invention relates to novel peptides, their use in treatment of chronic destruction of articular cartilage in autoimmune diseases, pharmaceutical compositions comprising said peptide, and a diagnostic method for the detection of autoreactive T cells in a test sample.

10       The immune system is established on a principle of discrimination between foreign antigens (non-self antigens) and autoantigens (self antigens, derived from the individuals own body) achieved by a build-in tolerance against the autoantigens.

      The immune system protects individuals against foreign antigens and responds to exposure to a foreign antigen by activating specific cells such as T- and B lymphocytes  
15       and producing soluble factors like interleukins, antibodies and complement factors. The antigen to which the immune system responds is degraded by the antigen presenting cells (APCs) and a fragment of the antigen is expressed on the cell surface associated with a major histocompatibility complex (MHC) class II glycoprotein. The MHC-glycoprotein-antigen-fragment complex is presented to a T cell which by virtue of its T  
20       cell receptor recognizes the antigen fragment conjointly with the MHC class II protein to which it is bound. The T cell becomes activated, i.e. proliferates and/or produces interleukins, resulting in the expansion of the activated lymphocytes directed to the antigen under attack (Grey et al., Sci. Am., 261:38-46, 1989).

      Self antigens are also continuously processed and presented as antigen fragments by  
25       the MHC glycoproteins to T cells (Jardetsky et al., Nature 353:326-329, 1991). Self recognition thus is intrinsic to the immune system. Under normal circumstances the immune system is tolerant to self antigens and activation of the immune response by these self antigens is avoided.

When tolerance to self antigens is lost, the immune system may become activated against one or more self antigens, resulting in the activation of autoreactive T cells and the production of autoantibodies. This phenomenon is referred to as autoimmunity. As the immune response in general is destructive, i.e. meant to destroy the invasive foreign antigen, autoimmune responses can cause destruction of the body's own tissue.

The contribution of T cells to autoimmune diseases has been established in several studies. In mice, experimental autoimmune encephalomyelitis (EAE) is mediated by a highly restricted group of T cells, linked by their specificity for a single epitope of myelin basic protein (MBP) complexed to an MHC class II molecule. In the Lewis rat, a species with high susceptibility to various autoimmune diseases, disease has been shown to be mediated by T cells. In humans autoimmune diseases are also thought to be associated with the development of auto-aggressive T cells.

A destructive autoimmune response has been implicated in various diseases such as rheumatoid arthritis (RA), in which the integrity of articular cartilage is destroyed by a chronic inflammatory process resulting from the presence of large numbers of activated lymphocytes and MHC class II expressing cells. The mere presence of cartilage appears necessary for sustaining the local inflammatory response: it has been suggested that cartilage degradation is associated with the activity of cartilage-responsive autoreactive T cells in RA (Sigall et al., Clin. Exp. Rheumat. 6:59, 1988; Glant et al., Biochem. Soc. Trans. 18:796, 1990; Burmester et al., Rheumatoid arthritis Smolen, Kalden, Maini (Eds) Springer-Verlag Berlin Heidelberg, 1992). Furthermore, removal of cartilage from RA patients by surgery was shown to reduce the inflammatory process (R.S. Laskin, J. Bone Joint Surgery (Am) 72:529, 1990). The cartilage proteins are therefore considered to be target autoantigens which are competent of stimulating T cells. Activation of these autoreactive T cells leads to development of autoimmune disease. However, the identification of the autoantigenic components that play a role in the onset of rheumatoid arthritis has so far remained elusive.

30

The inflammatory response resulting in the destruction of the cartilage can be treated by several drugs, such as for example steroid drugs. However, these drugs are often immunosuppressive drugs that are nonspecific and have toxic side effects. The

disadvantages of nonspecific immunosuppression makes this a highly unfavourable therapy.

The antigen-specific, nontoxic immunosuppression therapy provides a very attractive alternative for the nonspecific immunosuppression. This antigen-specific therapy involves the treatment of patients with the target autoantigen or with synthetic T cell-reactive peptides derived from the autoantigen. These synthetic peptides correspond to T cell epitopes of the autoantigen and can be used to induce specific T cell tolerance both to themselves and to the autoantigen. Although it seems paradoxical to desensitize the immune system with the very same antigen responsible for activating the immune system, the controlled administration of the target (auto)antigen can be very effective in desensitization of the immune system. Desensitization or immunological tolerance of the immune system is based on the long-observed phenomenon that animals which have been fed or have inhaled an antigen or epitope are less capable of developing a systemic immune response towards said antigen or epitope when said antigen or epitope is introduced via a systemic route.

The human cartilage glycoprotein-39 (HC gp-39) was previously identified as a target autoantigen in rheumatoid arthritis (RA) (Verheijden et al., *Arthritis Rheum.* 40:1115-1125, 1997). The strategy followed for identification of relevant auto-epitopes within HC gp-39 was based on the assumption that the DR4 or DR1 molecules predispose to RA (Gao et al., *Arthritis Rheum.* 33:939-946, 1990; Nelson et al., *Rheumatoid Arthritis*, In Proceedings of the Eleventh International Histocompatibility Workshop and Conference. Vol 1, Tsuji et al Ed, Oxford University Press, 1991) at two levels, firstly, by shaping the T cell repertoire and secondly, by determinant selection. The shared epitope found among the RA-associated DR molecules might be involved in selection of similar sets of peptides for presentation to T cells (Gregerson et al., *Arthritis Rheum.* 30:1205-1213, 1987). Putative binding sequences within the primary structure of HC gp-39 were identified by use of a DR4 (B1\*0401) peptide binding motif (Verheijden et al., *Arthritis Rheum.* 40:1115-1125, 1997). HC gp-39, a protein of 362 amino acids, excluding the signal sequence (Hakala et al., *J. Biol. Chem.* 268:25803-25810, 1993), contains six regions accommodating this motif. Four peptides thus selected were synthesized and tested for binding the RA-associated DR1 and DR4 (B1\*0401 and 0404) variants. All motif-based peptides, spanning residues 103-116, 259-275, 263-275 and 326-338 of HC gp-39, were found to bind with high relative affinity to DRB1\*0401 molecules. The recognition of these peptides by peripheral blood T cells from RA patients and healthy donors was subsequently examined. All motif-

based peptides were readily recognized in RA patients, thereby suggesting a high frequency of HC gp-39-specific T cells in RA. The response to 263-275 was most prominent; 8 out of 18 RA patients responded to this peptide (Verheijden et al., Arthritis Rheum. 40:1115-1125, 1997). Thus, HC gp-39 is a target for immune recognition in the joint.

The significance of this protein for arthritic disease was further demonstrated by its arthritogenicity in Balb/c mice. A single injection in the chest region with  $\mu$ g amounts of protein mixed in IFA, induced a chronic joint inflammation reminiscent of RA (Verheijden et al., Arthritis Rheum. 40:1115-1125, 1997).

Recently, a novel human chondrocyte protein, YKL-39, was isolated and described (Hu et al., J.Biol.Chem. 271: 19415-19420, 1996). The protein shares significant sequence identity with HC gp-39 (YKL-40). Another homologue of HC gp-39 is secreted by human macrophages and is termed chitotriosidase (Boot et al., J.Biol.Chem. 270: 26252-26256, 1995). The sequences corresponding to the HC gp-39 (263-275) peptide RSFTLASSETGVG (SEQ ID NO:3) are identified as HSFTLASAETTVG (SEQ ID NO:2) within the YKL-39 protein (266-278) and as RSFTLASSSDTRVG (SEQ ID NO:4) within macrophage chitotriosidase (269-282) respectively (Table 1).

The chitotriosidase peptide Chi (269-282) contains the DRB1\*0401 peptide binding motif which was previously used for selection of T-cell epitopes within proteins. In contrast, the YKL-39 (266-278) peptide does not contain this 0401 motif.

It will be clear that tolerization of HC gp-39 (263-275)-reactive T-cells may be of benefit to RA patients. Likewise, mimicry epitopes of HC gp-39 (263-275) may have a similar function and may be used to induce tolerance. Preferably such mimicry epitopes will have at least the same tolerizing capacity.

To effectively use tolerance induction therapy to treat T cell mediated cartilage destruction, there is a great need to identify T cell-reactive peptides which can desensitize patients against the autoantigen that is activating the T cells responsible for the inflammatory process.

Although the YKL-39 peptide does not contain the 0401 motif, it was surprisingly found that the YKL-39 (266-278) epitope is a mimicry epitope of HC gp-39 (263-275).

This epitope therefore is useful for tolerization of autoreactive T-cells with reactivity to HC gp-39 (263-275), YKL-39 (266-278) or their mimicry epitopes in rheumatoid arthritis patients.

5 It is an object of the invention to provide peptides which are able to induce systemic immunological tolerance, more in particular specific T cell tolerance, preferably to the responsible cartilage antigen in patients suffering from T cell-mediated cartilage destruction. The peptides of the present invention are characterized in that they comprise one or more of the amino acid sequences FTLASAETT (SEQ ID NO: 1). More  
10 specifically, a peptide according to the invention comprises HSFTLASAETTVG (SEQ ID NO: 2).

Also within the scope of the invention are multimers of the peptides according to the invention such as for example a dimer or trimer of the peptides according to the invention. A multimer according to the invention can either be a homomer, consisting of  
15 a multitude of the same peptide, or a heteromer consisting of different peptides.

The characteristic amino acid sequences of the peptides according to the invention can be flanked by random amino acid sequences. Preferred are flanking sequences, that have a stabilizing effect on the peptides, thus increasing their biological availability.

Human Cartilage glycoprotein 39 is a target autoantigen in RA patients which  
20 activates specific T cells, thus causing or mediating the inflammatory process. HC gp-39 derived peptides were predominantly recognized by autoreactive T cells from RA patients but rarely by T cells from healthy donors, thus indicating that HC gp-39 is an autoantigen in RA. The arthritogenic nature of HC gp-39 was further substantiated in the Balb/c mouse. A single, subcutaneous injection of said protein in Balb/c mice was able  
25 to initiate arthritic signs in the animals. The course of the HC gp-39- induced disease was characterized by relapses occurring periodically in fore paws and/or hind paws and gradually developed from a mild arthritis into a more severe form. Also, a symmetrical distribution of afflicted joints was observed which is, together with the observation of recurrent relapses, reminiscent of disease progression in arthritis, especially RA.

30 It was surprisingly found that the YKL-39 266-278 peptide was effective as a tolerogen. It will be clear to those skilled in the art that the peptides may be extended at either side of the peptide or at both sides and still exert the same immunological function. The extended part may be an amino acid sequence similar to the natural sequence of the protein YKL-39

The peptides according to the invention can be prepared by well known organic chemical methods for peptide synthesis such as, for example, solid-phase peptide synthesis described for instance in J. Amer. Chem. Soc. 85:2149 (1963) and Int. J. Peptide Protein Res. 35:161-214 (1990). The peptides according to the invention can also  
5 be prepared by recombinant DNA techniques. A nucleic acid sequence coding for a peptide according to the invention or a multimer of said peptides is inserted into an expression vector. Suitable expression vectors comprise the necessary control regions for replication and expression. The expression vector can be brought to expression in a host cell. Suitable host cells are, for instance, bacteria, yeast cells and mammalian cells. Such techniques are  
10 well known in the art, see for instance Sambrooke et al, Molecular Cloning: a Laboratory Manual, Cold Spring Harbor laboratory Press, Cold Spring Harbor, 1989.

The peptides may be stabilised by C- and/or N- terminal modifications, which will decrease exopeptidase catalysed hydrolysis. The modifications may include: C-terminal  
15 acylation, (e.g. acetylation = Ac-peptide), N-terminal amide introduction, (e.g. peptide-NH<sub>2</sub>) combinations of acylation and amide introduction (e.g. Ac-peptide-NH<sub>2</sub>) and introduction of D-amino acids instead of L-amino acids (Powell et al., J. Pharm. Sci., 81:731-735, 1992).

Other modifications are focussed on the prevention of hydrolysis by  
20 endopeptidases. Examples of these modifications are: introduction of D-amino acids instead of L-amino acids, modified amino acids, cyclisation within the peptide, introduction of modified peptide bonds, e.g. reduced peptide bonds  $\psi[\text{CH}_2\text{NH}]$  and e.g. peptoids (N-alkylated glycine derivatives) (Adang et al, Recl. Trav. Chim. Pays-Bas, 113:63-78, 1994 and Simon et al, Proc. Natl. Acad. Sci. USA, 89:9367-9371, 1992).

25

The peptides according to the invention are T-cell epitopes, which are recognized by and are able to stimulate autoreactive T-cells. These autoreactive T cells may be found e.g. in the blood of patients suffering from autoimmune diseases.

Thus, according to the invention the peptides, said peptides resembling the MHC  
30 Class II restricted T-cell epitopes present on the target autoantigen comprising the peptide of SEQ ID NO:1 or SEQ ID NO:2, are very suitable for use in a therapy to induce specific T-cell tolerance to said autoantigen in mammals, more specifically humans, suffering from T-cell mediated cartilage destruction, such as for example arthritis. more specifically rheumatoid arthritis. Optionally such a treatment can be



combined with the the administration of other medicaments such as DMARDs (Disease Modifying Anti-Rheumatic Drugs e.g. sulfasalazine, anti-malarials (chloroquine, hydroxychloroquine) injectable or oral gold, methotrexate, D-penicillamine, azathioprine, cyclosporine, mycophenolate), NSAIDs (non steroidal anti inflammatory drugs), corticosteroids or other drugs knowns to influence the course of the disease in  
5 autoimmune patients.

The peptides according to the invention can also be used to modulate lymphocytes that are reactive to antigens other than said autoantigen but are present in the same tissue  
10 as the autoantigen i.e. proteins or parts thereof comprising the peptide according to SEQ ID NO:1 or SEQ ID NO:2. By the induction of antigen-specific T-cell tolerance, autoimmune disorders can be treated by bystander suppression. More in general, the cells to be modulated are hematopoietic cells. In general, in order to function as a tolerogen the peptide must fulfill at least two conditions i.e. it must possess an immune  
15 modulating capacity and it must be expressed locally usually as part of a larger protein.

Thus, the present invention provides a method to treat patients suffering from inflammatory autoimmune diseases, by administration of a pharmaceutical preparation comprising the peptide according to the invention. Such patients may suffer from diseases like Graves' diseases, juvenile arthritis, primary glomerulonephritis, osteoarthritis, Sjögren's syndrome, myasthenia gravis, rheumatoid arthritis, Addison's  
20 disease, primary biliary sclerosis, uveitis, systemic lupus erythematosus, inflammatory bowel disease, multiple sclerosis or diabetes. The peptides according to the present invention therefore can be used in the preparation of a pharmaceutical to induce tolerance in patients suffering from these diseases.

25 Treatment of autoimmune disorders with the peptides according to the invention makes use of the fact that bystander suppression is induced to unrelated but co-localized antigens. The regulatory cells secrete in an antigen specific fashion pleiotropic proteins such as cytokines which may downmodulate the immune response.

30 According to the invention, patients suffering from T-cell mediated destruction of the articular cartilage can be treated with a therapeutical composition comprising one or more peptides according to the invention and a pharmaceutical acceptable carrier. Administration of the pharmaceutical composition according to the invention will

autoreactive T cells of these patients, to the autoantigenic proteins in the articular cartilage under attack and other self antigens which display the identified MHC Class II binding T cell epitopes characterized or mimicked by the amino acid sequences of one or more of the peptides according to the invention. The induced tolerance thus will lead to a reduction of the local inflammatory response in the articular cartilage under attack.

Very suitable peptides to be used in a pharmaceutical composition according to the invention are the peptides comprising the YKL-39 (268-276) or the YKL-39 (266-278) peptide flanked by sequences up to a total length of 55 amino acids. More preferably the peptides have a length of 25 amino acids. Even more preferably the amino acid sequence of the peptides is FTLASAETT or HSFTLASAETTVG.

The peptides according to the invention have the advantage that they have a specific effect on the autoreactive T cells thus leaving the other components of the immune system intact as compared to the nonspecific suppressive effect of immunosuppressive drugs. Treatment with the peptides according to the invention will be safe and no toxic side effects will occur.

Systemic immunological tolerance can be attained by administering high or low doses of peptides according to the invention. The amount of peptide will depend on the route of administration, the time of administration, the age of the patient as well as general health conditions and diet.

In general, a dosage of 0.01 to 10000 µg of peptide per kg body weight, preferably 0.05 to 500 µg, more preferably 0.1 to 100 µg of peptide can be used.

Pharmaceutical acceptable carriers are well known to those skilled in the art and include, for example, sterile saline, lactose, sucrose, calcium phosphate, gelatin, dextrin, agar, pectin, peanut oil, olive oil, sesame oil and water. Other carriers may be, for example MHC class II molecules, if desired embedded in liposomes.

In addition the pharmaceutical composition according to the invention may comprise one or more adjuvants. Suitable adjuvants include, amongst others, aluminium hydroxide, aluminium phosphate, amphigen, tocophenols, monophosphenyl lipid A, muramyl dipeptide and saponins such as Quill A. Preferably, the adjuvants to be used in the tolerance therapy according to the invention are mucosal adjuvants such as the cholera toxine B-subunit or carbomers, which bind to the mucosal epithelium. The amount of adjuvant depends on the nature of the adjuvant itself.

Furthermore the pharmaceutical composition according to the invention may comprise one or more stabilizers such as, for example, carbohydrates including sorbitol, mannitol, starch, sucrosextrin and glucose, proteins such as albumin or casein, and buffers like alkaline phosphates.

- 5        Suitable administration routes are e.g. intramuscular injections, subcutaneous injections, intravenous injections or intraperitoneal injections, oral administration and nasal administration such as sprays.

10        It is another object of the invention to provide a method for detecting autoreactive T cells involved in the destruction of articular cartilage and test kits to be used in said method. Thus, the peptides according to the invention are also very suitable for use in a diagnostic method to detect the presence of activated autoreactive T cells involved in the chronic inflammation and destruction of the articular cartilage.

The diagnostic method according to the invention comprises the following steps:

- 15        a) isolation of the peripheral blood mononuclear cells (PBMC) from a blood sample of an individual,
- b) culture said PBMC under suitable conditions,
- c) incubation of said PBMC culture in the presence of one or more peptides according to the invention, and
- 20        d) detection of a response of T cells, for example a proliferative response, indicating the presence of activated autoreactive T cells in the individual.

The detection of a proliferative response of T cells can be detected by, for example, the incorporation of  $^3\text{H}$ -thymidine.

25        Also within the scope of the invention are test kits which comprise one or more peptides according to the invention. These test kits are suitable for use in a diagnostic method according to the invention.

The following examples are illustrative for the invention and should in no way be interpreted as limiting the scope of the invention.

## Legends to the figures

### Figure 1

Figure 1a, b, c. Cross reactivity of three, different, HC gp-39-specific hybridomas (8B12, 14G11, 20H5) with YKL-39 (266-278)

5 (CVR0271B = HC gp-39 (263-275), KV0432B = YKL-39 (266-278), CC0332B = Chi (269-282), KV0431A = YKL-39 (262-274). HCDA.8B12.1D8, 14G11.1H7 and 20H5.4F6.2F6 are HLA-DRB1\*0401-restricted hybridomas specific for HC gp-39 (263-275). Activation of T-cell hybridomas is expressed as IL-2 production.

### Figure 2 In vivo tolerization with HC gp-39 (263-275) or YKL-39 (266-278)

10 Balb/c mice were tolerized by intranasal application of 50, 10 or 2 microgram of HC gp-39 (263-275) or YKL-39 (266-278) followed by immunization with HC gp-39 (263-275). Mice that were pretreated with saline or that were left untreated were included as controls.

15

## Examples

### Example 1 Alignment of sequences

The human chondrocyte protein, YKL-39 shares significant sequence identity with HC gp-39 (YKL-40). Another homologue of HC gp-39 is secreted by human  
20 macrophages and is termed chitotriosidase (Boot et al., 1995). The sequences corresponding to RSFTLASSETGVG (HC gp-39 (263-275), SEQ ID NO:3) were identified as HSFTLASAETTVG within the YKL-39 protein (266-278) and as RSFTLASSSDTRVG (SEQ ID NO:4) within macrophage chitotriosidase (269-282) respectively (Table 1). Chi (269-282) contains the HLA-DRB1\*0401 peptide binding  
25 motif which was previously used for selection of T-cell epitopes within proteins. In contrast, the YKL-39 (266-278) peptide does not contain this motif. All peptides were synthesized.

Table 1. Alignment of the HC gp-39 (263-275) sequence with the corresponding region in YKL-39 and macrophage Chitotriosidase

HCgp-39 263-275	R	S	F	T	L	A	S	S	-	E	T	G	V	G
YKL-39 266-278	H	S	F	T	L	A	S	A	-	E	T	T	V	G
Chi (269-282)	R	S	F	T	L	A	S	S	S	D	T	R	V	G

### Example 2 Binding of peptides to HLA-DRB1\*0401

The peptides from example 1 were tested for binding the DRB1\*0401-encoded molecules. HLA-DR4 (DRB1\*0401) molecules were purified from the homozygous EBV-transformed human B lymphoblastoid cell lines Huly138IC2 and the competition peptide HLA-DR binding assay was performed basically as described by Verheijden et al., 1997. The affinity of a given peptide for binding DRB1\*0401-encoded molecules was related to competition with a marker peptide. This relative binding affinity was defined as the peptide concentration at which the signal was reduced to 50% (IC<sub>50</sub>). The HA-F peptide is a positive control (Hemagglutinin 307-319; PKFVKQNTLKLAT; at position 309 Y is substituted by F; SEQ ID NO:5). The peptide is known to have a high affinity for DRB1\*0401 molecules.

As expected, the Chi(269-282) peptide was found to bind with high affinity to DRB1\*0401 (see table 2). The YKL-39 (266-278) peptide, which does not accommodate the effective DRB1\*0401 peptide binding motif, bound with very high affinity to DR4 (B1\*0401).

Table 2 Peptide binding to HLA-DRB1\*0401-encoded molecules

peptide	batch	IC <sub>50</sub> values		
		Exp. A	Exp. B	Exp. C
YKL39(262-274)	KV0431A	0.006	0.005	ND
YKL39(266-278)	KV432B	0.035	0.032	0.12
HCgp39(263-275)	CVR271B	ND	0.008	0.038
Chi(269-282)	CC0332B	0.053	0.11	0.16
HA-F	AE0690A	0.20	0.14	0.20

### Example 3 Stimulation T-cell hybridomas

Hybridomas specific for HC gp-39 (263-275) were tested for recognition of the corresponding sequences.

To test the cross reactivity of the 3 different, HC gp-39-specific hybridoma cell lines with the YKL-39 or the chitotriosidase peptide,  $5 \times 10^4$  hybridoma cells and  $2 \times 10^5$  irradiated (12000 RAD), EBV-transformed B cells carrying the DRB1\*0401 specificity were incubated in 150  $\mu$ l volumes in wells of a round-bottomed microtiter plate. Peptide antigen (HC gp-39 (263-275), YKL-39 (266-278), chitotriosidase (269-282) or a control peptide) was added in 50  $\mu$ l volumes to duplicate wells. Forty-eight hr later 100  $\mu$ l of the culture supernatant was assayed for IL-2 production using a sandwich ELISA with Pharmingen antibodies specific for mouse IL-2.

It was found that the synthetic peptide YKL-39 (266-278) generated a response similar to HC gp-39 (263-275) whereas the Chi (269-282) did not generate a response. The data suggest that the three different TCRs utilized by three different hybridomas do not discriminate between HC gp-39 (263-275) or YKL-39 (266-278) when presented by DRB1\*0401-encoded molecules (Figure 1a, b, c) but do discriminate between HC gp-39 (263-275) and Chi (269-282). The data indicate that YKL-39 (266-278) is a mimicry epitope of HC gp-39 (263-275). (Fig 1a,b,c)

### Example 4 Recognition of YKL-39 (266-278) by PBMC

Peripheral blood mononuclear cells (PBMC) were isolated from heparinized peripheral blood by standard centrifugation on Ficoll-Paque (Pharmacia, Uppsala, Sweden). Cells were suspended in wells of a 24 well plate in a concentration of  $5 \times 10^5$  cells per ml. Cells were incubated in medium alone or in the presence of 10 or 50  $\mu$ g/ml peptide antigen (YKL-39 (266-278)). Cultures were incubated for 6 days at 37 °C in a humidified atmosphere of 5% CO<sub>2</sub>. Cells were then suspended and 100 or 150  $\mu$ l volumes of medium was distributed in 4-fold in wells of a 96-well roundbottomed plate. Cells were then pulsed with 0.5  $\mu$ Ci ( $1.85 \times 10^4$  Bq) [<sup>3</sup>H]thymidine ([<sup>3</sup>H]TdR) and 18 hr later incorporated radioactivity was measured. Results as shown in table 3 are expressed as stimulation indices (SI) (antigen-specific counts/background counts)

From Table 3a it can be concluded that the YKL-39 (266-278) epitope is readily recognized in RA patients. Table 3b indicates that recognition of YKL-39 (266-278) by

PBMC coincides with recognition of HC gp-39 (263-275) and HC gp-39 and furthermore that recognition of YKL-39 (266-278) is generally more pronounced than recognition of HC gp-39 (263-275).

5 Table 3a. Recognition of the YKL-39 (266-278) epitope by PBMC from RA patients.

Donor		typing	SI	SI
			10 µg/ml	50 µg/ml
242-0.2	NR	0404/15	3	<2
337-0.2	R	0401/02	19	58
338-0.1	NR	03/14	<2	<2
454-0	R	0401/	9	9
456-0	R	ND	15	4
457-0	NR	ND	<2	<2
458-0	R	ND	4	27
459-0	R	ND	<2	25
460-0	NR	ND	3	<2

*SI = antigen-specific counts/background counts. SI ≥ 5 are regarded positive R = responder, NR = non-responder*

Table 3b. Recognition of YKL-39 (266-278) coincides with recognition of HC gp-39(263-275) and HC gp-39 protein.

Donor	R/N	YKL-39(266-278)		HC gp-39(263-275)		HC gp-39	
		SI	SI	SI	SI	SI	SI
	$\mu\text{g/ml}$	10	50	10	50	10	50
169	R	27	32	10	27	24	44
455	R	20	35	1	15	45	95
447	NR	1	2	1	1	1	1
327	R	6	5	3	5	12	19

SI = antigen-specific counts/background counts. SI  $\geq$  5 are regarded positive. R = responder, NR = non-responder. NT = not tested. Donor 447 responds to Tetanus toxoid and Candida Albicans.

#### Example 5 Tolerance induction

A HC gp-39 (263-275)-specific DTH assay suitable to monitor tolerance induction with peptide antigens was developed. Immunisation of Balb/c mice with HC gp-39 (263-275) in incomplete Freund's adjuvant (IFA) was found to be effective in the induction of a DTH response following challenge with the HC gp-39 (263-275) peptide. This peptide-based DTH system was used to detect modulation of the DTH response by nasal application of HC gp-39 (263-275) peptide. It was found that nasal application of HC gp-39 (263-275), in a dose-dependent manner, downmodulated the HC gp-39 (263-275)-induced DTH response. Nasal application of YKL-39 (266-278), however, resulted in a more enhanced downmodulation of the DTH response, indicating that YKL-39 (266-278) can efficiently tolerate a peptide-specific response induced with HC gp-39 (263-275) (Table 4, Figure 2a,b,c).



Table 4. Experimental set-up tolerization experiment

Pretreatment	sensibilisation	challenge	tolerance
none	HC gp-39 (263-275)	HC gp-39 (263-275)	no
saline	HC gp-39 (263-275)	HC gp-39 (263-275)	no
HC gp-39 (263-275)	HC gp-39 (263-275)	HC gp-39 (263-275)	yes
YKL-39 (266-278)	HC gp-39 (263-275)	HC gp-39 (263-275)	yes


## Claims

1. Peptide having an amino acid sequence of 9-55 amino acid residues comprising the amino acid sequence FTLASAETT (SEQ ID NO:1).
- 5 2. Peptide according to claim 1 comprising the amino acid sequence HSFTLASAETTVG (SEQ ID NO:2).
3. Peptide according to claim 1 or 2 having an amino acid sequence of up to 25 amino acid residues.
4. Peptide according to claim 1 or 2 having the amino acid sequence FTLASAETT (SEQ ID NO:1) or HSFTLASAETTVG (SEQ ID NO:2).
- 10 5. Peptides according to any of the claims 1-4 for use as a therapeutic substance.
6. Pharmaceutical composition comprising one or more of the peptides according to claims 1-4, and a pharmaceutical acceptable carrier.
7. Use of one or more of the peptides according to claims 1-4 for the manufacture of a pharmaceutical preparation for the induction of specific T-cell tolerance to an autoantigen in patients suffering from autoimmune disorders, more specifically arthritis.
- 15 8. Diagnostic composition comprising one or more of the peptides according to any of the claims 1-4 and a detection agent.

1/6

Figure 1A


Clone 8B12.1D8



2/6

Figure 1B


Clone 14G11.1H7



3/6

Figure 1C.

Clone 20H5.4F6.2F6



4/6




Figure 2

5/6

challenge peptide:  
HC gp-39 (263-275)




Figure 2, cont.

6/6

challenge peptide:  
HC gp-39 (263-275)




Figure 2, cont.



## SEQUENCE LISTING

<110> Akzo Nobel N.V.

5 <120> Novel peptides for use in immunotherapy of autoimmune diseases

<130>

10 <140>  
<141>

<150> 98202470.5

15 <151> 1998-07-23

<160> 5

<170> PatentIn Ver. 2.1

20 <210> 1  
<211> 9  
<212> PRT  
<213> Homo sapiens

25 <400> 1  
Phe Thr Leu Ala Ser Ala Glu Thr Thr  
1 5

30 <210> 2  
<211> 13  
<212> PRT

35 <213> Homo sapiens

<400> 2  
His Ser Phe Thr Leu Ala Ser Ala Glu Thr Thr Val Gly  
1 5 10

40 <210> 3  
<211> 13  
<212> PRT

<213> Homo sapiens

<400> 3

Arg Ser Phe Thr Leu Ala Ser Ser Glu Thr Gly Val Gly

5

1

5

10

<210> 4

10

<211> 14

<212> PRT

<213> Homo sapiens

<400> 4

15

Arg Ser Phe Thr Leu Ala Ser Ser Ser Asp Thr Arg Val Gly

1

5

10

20

<210> 5

<211> 13

<212> PRT

<213> Homo sapiens

25

<400> 5

Pro Lys Phe Val Lys Gln Asn Thr Leu Lys Leu Ala Thr

1

5

10

30

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C07K 14/47, G01N 33/68, A61K 38/08,</b> <b>38/10, C07K 7/06, 7/08</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 00/05254</b> <b>(43) International Publication Date:</b> 3 February 2000 (03.02.00)
<b>(21) International Application Number:</b> PCT/EP99/05050 <b>(22) International Filing Date:</b> 16 July 1999 (16.07.99) <b>(30) Priority Data:</b> 98202470.5                      23 July 1998 (23.07.98)                      EP <b>(71) Applicant (for all designated States except US):</b> AKZO NOBEL N.V. [NL/NL]; Velperweg 76, NL-6824 BM Arnhem (NL). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> VERHEIJDEN, Gijsbertus, Franciscus, Maria [NL/NL]; Heischouw 7, NL-5345 XT Oss (NL). BOOTS, Anna, Maria, Helena [NL/NL]; Verlengde Torenstraat 10, NL-5366 AV Megen (NL). <b>(74) Agent:</b> HERMANS, F., G., M.; P.O. Box 20, NL-5340 BH Oss (NL).	<b>(81) Designated States:</b> AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>  <b>(88) Date of publication of the international search report:</b> 15 June 2000 (15.06.00)	

**(54) Title:** NOVEL PEPTIDES FOR USE IN IMMUNOTHERAPY OF AUTOIMMUNE DISEASES**(57) Abstract**

The invention relates to the use of novel peptides in a peptide induced tolerance therapy to prevent autoimmune disorders and in particular their use in treatment of chronic destruction of articular cartilage. The invention furthermore embraces pharmaceutical compositions comprising said peptides and a diagnostic method for the detection of autoreactive T cells in a test sample.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark			SE	Sweden		

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 99/05050

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07K14/47 G01N33/68 A61K38/08 A61K38/10 C07K7/06  
C07K7/08

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K A61K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 06859 A (HUMAN GENOME SCIENCES INC ; THOTAKURA NAGESWARA RAO (US); CHOPRA AR) 19 February 1998 (1998-02-19) Residues 334-342 of hpmfw51 figure 3 ---	
A	EP 0 805 206 A (SMITHKLINE BEECHAM CORP) 5 November 1997 (1997-11-05) reidues 306-314 of seq ID 2. page 3, last paragraph ---	
A	WO 96 13517 A (AKZO NOBEL NV ; BOOTS ANNA MARIA HELENA (NL); VERHEIJDEN GIJSBERTUS) 9 May 1996 (1996-05-09) abstract --- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

21 March 2000

Date of mailing of the international search report

28/03/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL 2000 LV Dordrecht

Authorized officer

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/05050

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WO 97 40068 A (AKZO NOBEL NV ;BOOTS ANNA MARIA HELENA (NL); VERHEIJDEN GIJSBERTUS)  30 October 1997 (1997-10-30)  seq id 41</p> <p>-----</p>	

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/05050

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9806859 A	19-02-1998	AU 6693996 A	06-03-1998
EP 0805206 A	05-11-1997	JP 10066585 A	10-03-1998
WO 9613517 A	09-05-1996	AT 177756 T	15-04-1999
		AU 696827 B	17-09-1998
		AU 3925295 A	23-05-1996
		BR 9506377 A	16-09-1997
		CN 1168677 A	24-12-1997
		DE 69508380 D	22-04-1999
		DE 69508380 T	09-09-1999
		EP 0733065 A	25-09-1996
		ES 2130672 T	01-07-1999
		FI 962619 A	25-06-1996
		GR 3030296 T	30-09-1999
		HU 74847 A	28-02-1997
		JP 9507861 T	12-08-1997
		NO 962695 A	26-06-1996
		NZ 295659 A	26-02-1998
		PL 315198 A	14-10-1996
		TR 960361 A	21-06-1996
		US 5736507 A	07-04-1998
WO 9740068 A	30-10-1997	AU 2768597 A	12-11-1997
		BR 9708744 A	03-08-1999
		CA 2251680 A	30-10-1997
		CN 1216551 A	12-05-1999
		CZ 9803406 A	17-02-1999
		EP 0920451 A	09-06-1999
		HU 9901375 A	30-08-1999
		NO 984937 A	01-12-1998
		PL 329618 A	29-03-1999
		ZA 9703071 A	05-08-1998

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**